ELSEVIER

Environmental Pollution 154 (2008) 504–507

ENVIRONMENTAL POLLUTION

ww.elsevier.com/locate/envpol

Contents of Volume 154

Number 1 2008

SPECIAL ISSUE: Mercury Cycling and Bioaccumulation in the Environment

Guest editorial

1 **Bridging the knowledge gaps on the sources, speciation, fate and bioaccumulation of mercury in aquatic and terrestrial environments**
R. Harris, K. Kidd, J. Shanley

3 **Characteristics of mercury speciation in Minnesota rivers and streams**
S.J. Balogh, E.B. Swain, Y.H. Nollet

Methylmercury and inorganic mercury concentrations in four Minnesota streams were characterized to determine controlling variables.

12 **Seasonal variation in dissolved gaseous mercury and total mercury concentrations in Juam Reservoir, Korea**
J.-S. Park, S. Oh, M.-Y. Shin, M.-K. Kim, S.-M. Yi, K.-D. Zoh

The first seasonal measurements of DGM and THg concentrations in a Korean reservoir.

21 **Environmental mercury contamination of an artisanal zinc smelting area in Weining County, Guizhou, China**
G. Li, X. Feng, G. Qiu, X. Bi, Z. Li, C. Zhang, D. Wang, L. Shang, Y. Guo

Artisanal zinc smelting caused serious mercury contamination in Weining, Guizhou, China.

32 **Mercury cycling in surface water, pore water and sediments of Mugu Lagoon, CA, USA**
S.E. Rothenberg, R.F. Ambrose, J.A. Jay

Trends in mercury cycling differed between habitats and within matrices at Mugu Lagoon.

46 **Wetland influence on mercury fate and transport in a temperate forested watershed**
P. Selvendiran, C.T. Driscoll, J.T. Bushey, M.R. Montesdeoca

Wetlands are sources of THg and MeHg; the production of MeHg is seasonally dependent and driven by sulfate reduction in wetlands.

56 **The impact of eutrophication on the biogeochemical cycling of mercury species in a reservoir: A case study from Hongfeng Reservoir, Guizhou, China**
T. He, X. Feng, Y. Guo, G. Qiu, Z. Li, L. Liang, J. Lu

Eutrophication may affect the biogeochemical cycling of mercury in reservoir.

68 **Methylmercury in a predatory fish (*Cichla* spp.) inhabiting the Brazilian Amazon**
H. do A. Kehrig, B.M. Howard, O. Malm

Cichla spp. may be considered good bioindicators of methylmercury contamination in the Amazonian ecosystem because of their integration of this pollutant over time.

77 **Temporal changes in the distribution, methylation, and bioaccumulation of newly deposited mercury in an aquatic ecosystem**
D.M. Orihel, M.J. Paterson, P.J. Blanchfield, R.A. (Drew) Bodaly, C.C. Gilmour, H. Hintelmann

Mercury deposited to aquatic ecosystems becomes less available for uptake by biota over time.

89 **Correlates of mercury in fish from lakes near Clyde Forks, Ontario, Canada**
A.L.M. Ethier, A.M. Scheuhammer, D.E. Bond

Fish from lakes near a localized geological Hg source do not have elevated Hg concentrations.

98 **Total mercury, organic mercury and selenium in liver and kidney of a South American coastal dolphin**
T.G. Seixas, H. do A. Kehrig, M. Costa, G. Fillmann, A.P.M. Di Beneditto, E.R. Secchi, C.M.M. Souza, O. Malm, I. Moreira

This study presents some useful data on the bioaccumulation of Hg, organic Hg and Se, and their inter-element relationships in a Brazilian small cetacean.

107 **Lake variability: Key factors controlling mercury concentrations in New York State fish**
H.A. Simonin, J.J. Loukmas, L.C. Skinner, K.M. Roy

Lake water chemistry variables, dams, and wetlands play major roles in determining fish mercury concentrations.

116 **Influence of natural dissolved organic carbon on the bioavailability of mercury to a freshwater alga**
P.R. Gorski, D.E. Armstrong, J.P. Hurley, D.P. Krabbenhoft

Bioavailability of mercury to an alga was greatest at low concentrations of natural dissolved organic carbon and inhibited at high concentrations of natural dissolved organic carbon.

doi:10.1016/S0269-7491(08)00321-7

124 Wetlands as principal zones of methylmercury production in southern Louisiana and the Gulf of Mexico region
B.D. Hall, G.R. Aiken, D.P. Krabbenhoft, M. Marvin-DiPasquale, C.M. Swarzenski

This study, which presents the results of a landscape scale examination of methylmercury and total mercury cycling in southern Louisiana, shows that freshwater, brackish, and marine wetlands are important sites of methylmercury production, which could lead to increased fish Hg concentrations in the Gulf of Mexico region.

135 Characterization of heavy metal concentrations in the sediments of three freshwater rivers in Huludao City, Northeast China
N. Zheng, Q. Wang, Z. Liang, D. Zheng

Sediment in Wuli River, Cishan River, and Lianshan River has been contaminated by heavy metals and adverse effects would be expected frequently in Wuli River and Cishan River.

143 Comparison of total mercury and methylmercury cycling at five sites using the small watershed approach
J.B. Shanley, M. Alisa Mast, D.H. Campbell, G.R. Aiken, D.P. Krabbenhoft, R.J. Hunt, J.F. Walker, P.F. Schuster, A. Chalmers, B.T. Aulenbach, N.E. Peters, M. Marvin-DiPasquale, D.W. Clow, M.M. Shafer

High-flow sampling reveals strong contrasts in total mercury and methylmercury cycling in five diverse USA watersheds.

Number 2 2008

Review

155 Microbial degradation of the benzonitrile herbicides dichlobenil, bromoxynil and ioxynil in soil and subsurface environments – Insights into degradation pathways, persistent metabolites and involved degrader organisms
M.S. Holtze, S.R. Sørensen, J. Sørensen, J. Aamand

The benzonitrile herbicides dichlobenil, bromoxynil and ioxynil are liable to degrade to persistent metabolites posing a potential water contaminant problem.

Commentary

169 Exposure to inorganic arsenic from rice: A global health issue?
Y.-G. Zhu, P.N. Williams, A.A. Meharg

Regular papers

172 Organohalogen pollutants in herring from the northern Baltic Sea: Concentrations, congener profiles and explanatory factors
J. Koistinen, H. Kiviranta, P. Ruokojärvi, R. Parmanne, M. Verta, A. Hallikainen, T. Vartiainen

Sources of pollutants are reflected in levels and congener profiles of pollutants in Baltic herring.

184 A multi-stage sampling strategy for the delineation of soil pollution in a contaminated brownfield
S. Verstraete, M. Van Meirvenne

A multi-stage sampling strategy based on geostatistics provides an efficient procedure to delineate a pollution in a contaminated brownfield.

192 Monitoring aromatic hydrocarbon biodegradation by functional marker genes
M. Nyyssönen, R. Piskonen, M. Itävaara

Functional gene array analysis coupled with one-colour hybridization and internal standard based signal normalization provides efficient tool for monitoring contaminant biodegradation processes.

203 Toxic effects of arsenic on *Sinorhizobium–Medicago sativa* symbiotic interaction
E. Pajuelo, I.D. Rodríguez-Llorente, M. Dary, A.J. Palomares

First steps of nodulation of alfalfa, in particular infection thread formation, are more sensitive to As than nitrogen fixation due to plant effects.

212 Phytoextraction by arsenic hyperaccumulator *Pteris vittata* L. from six arsenic-contaminated soils: Repeated harvests and arsenic redistribution
M.I.S. Gonzaga, J.A.G. Santos, L.Q. Ma

Pteris vittata was effective in continuously removing arsenic from contaminated soils after three repeated harvests.

219 Developing climatic scenarios for pesticide fate modelling in Europe
S. Blenkinsop, H.J. Fowler, I.G. Dubus, B.T. Nolan, J.M. Hollis

The FOOTPRINT climatic zones provide an objective climatic classification and daily climate series that may be used for the modelling of pesticide fate across Europe.

232 Effects of broadleaf woodland cover on streamwater chemistry and risk assessments of streamwater acidification in acid-sensitive catchments in the UK
Z. Gagkas, K.V. Heal, N. Stuart, T.R. Nisbet

Broadleaf woodland <30% of catchment area should not pose a risk of increased streamwater nitrate and aluminium concentrations within acid-sensitive catchments.

241 Seasonal differences and within-canopy variations of antioxidants in mature spruce (*Picea abies*) trees under elevated ozone in a free-air exposure system
N. Hofer, M. Alexou, C. Heerdt, M. Löw, H. Werner, R. Matyssek, H. Rennenberg, K. Haberer

Antioxidant defence in sun and shade needles of Picea abies under free-air ozone fumigation in the seasonal course of two consecutive years.

254 Heavy metal (Cu, Zn, Cd and Pb) contamination of vegetables in urban India: A case study in Varanasi
R.K. Sharma, M. Agrawal, F.M. Marshall

Atmospheric depositions can significantly increase the heavy metal concentrations in vegetables during marketing.

264 Phytotoxicities of fluoranthene and phenanthrene deposited on needle surfaces of the evergreen conifer, Japanese red pine (*Pinus densiflora* Sieb. et Zucc.)
I. Oguntimehin, N. Nakatani, H. Sakugawa

Fluoranthene and phenanthrene caused negative effects on the needles of Japanese red pine.

272 Spatial trends of polybrominated diphenyl ethers in avian species: Utilization of stored samples in the Environmental Specimen Bank of Ehime University (es-Bank)
T. Kunisue, Y. Higaki, T. Isobe, S. Takahashi, A. Subramanian, S. Tanabe

Open sea birds have been exposed to PBDEs, but the accumulation levels were lower than those in Japanese coastal and inland birds.

283 Using trace element concentrations in *Corbicula fluminea* to identify potential sources of contamination in an urban river
G.L. Peltier, J.L. Meyer, C.H. Jagoe, W.A. Hopkins

C. fluminea collected downstream of CFPPs had elevated tissue concentrations of trace elements.

291 Microcosm investigations of stormwater pond sediment toxicity to embryonic and larval amphibians: Variation in sensitivity among species
J.W. Snodgrass, R.E. Casey, D. Joseph, J.A. Simon

Variation among species in sensitivity to pollutants can influence stormwater pond amphibian assemblages.

298 Distribution and sources of polycyclic aromatic hydrocarbons in surface sediments of rivers and an estuary in Shanghai, China
Y. Liu, L. Chen, Z. Jianfu, H. Qinghui, Z. Zhiliang, G. Hongwen

Surface sediments of two rivers and an estuary in Shanghai were contaminated by PAHs.

306 Timing of phosphate application affects arsenic phytoextraction by *Pteris vittata* L. of different ages
J.A.G. Santos, M.I.S. Gonzaga, L.Q. Ma, M. Srivastava

Young ferns coupled with split-P application were effective in arsenic removal by Pteris vittata.

312 Effect of field exposure to 38-year-old residual petroleum hydrocarbons on growth, condition index, and filtration rate of the ribbed mussel, *Geukensia demissa*
J.B. Culbertson, I. Valiela, Y.S. Olsen, C.M. Reddy

Mussels exposed to petroleum in transplant and field experiments exhibited significant long-term effects.

320 Phase partitioning modeling of ethanol, isopropanol, and methanol with BTEX compounds in water
K.Y. Lee

As a high-alcohol content fuel blend migrates through the subsurface and encounters water, phase partitioning of the mixture plays a critical role in the distribution and spreading of the pollutants.

330 Assessment of pesticide availability in soil fractions after the incorporation of winery-distillery vermicomposts
J.D. Fernández-Bayo, E. Romero, F. Schnitzler, P. Burauel

Winery vermicomposts as organic amendments to reduce pesticide pollution.

338 Copper desorption in flooded agricultural soils and toxicity to the Florida apple snail (*Pomacea paludosa*): Implications in Everglades restoration
T.C. Hoang, E.C. Rogevich, G.M. Rand, P.R. Gardinali, R.A. Frakes, T.A. Bargar

Copper desorbs from agricultural soils and is toxic to the Florida apple snail.

348 Linear alkylbenzenes in riverine runoff of the Pearl River Delta (China) and their application as anthropogenic molecular markers in coastal environments
H.-G. Ni, F.-H. Lu, J.-Z. Wang, Y.-F. Guan, X.-L. Luo, E.Y. Zeng

Occurrence of LABs in riverine runoff of the Pearl River Delta (China) was examined.

Number 3 2008

SPECIAL ISSUE: Reduced Nitrogen in Ecology and the Environment

Guest editorial

357 Reduced nitrogen in ecology and the environment: Special issue of the ESF-FWF Conference in partnership with LFUI, October 2006
J.W. Erisman, M.A. Sutton

359 Reduced nitrogen has a greater effect than oxidised nitrogen on dry heathland vegetation
L.J.L. van den Berg, C.J.H. Peters, M.R. Ashmore, J.G.M. Roelofs

Reduced nitrogen and high NH_4^+/NO_3^- ratios in deposition are responsible for a decline of forbs in Western European heathlands.

370 Modelling the spatial distribution of ammonia emissions in the UK
S. Hellsten, U. Dragosits, C.J. Place, M. Vieno, A.J. Dore, T.H. Misselbrook, Y.S. Tang, M.A. Sutton

It is important to provide robust estimates of the spatial distribution of ammonia emissions, since the model output is used to assess potential environmental impacts, e.g. through the exceedance of critical loads.

380 Causes of change in nitrophytic and oligotrophic lichen species in a Mediterranean climate: Impact of land cover and atmospheric pollutants
P. Pinho, S. Augusto, M.A. Martins-Loução, M.J. Pereira, A. Soares, C. Máguas, C. Branquinho

Causes of change in nitrophytic and oligotrophic lichen species.

Review

390 Relationship between ammonia stomatal compensation point and nitrogen metabolism in arable crops: Current status of knowledge and potential modelling approaches
R.S. Massad, B. Loubet, A. Tuzet, P. Cellier

A model for ammonia stomatal compensation point at the leaf level scale was developed.

404 Stress responses of *Calluna vulgaris* to reduced and oxidised N applied under 'real world conditions'
L.J. Sheppard, I.D. Leith, A. Crossley, N. Van Dijk, D. Fowler, M.A. Sutton, C. Woods

Exposure to NH_3 reduces stress resistance and increases visible damage in mature Calluna.

414 Heterogeneity of soil surface ammonium concentration and other characteristics, related to plant specific variability in a Mediterranean-type ecosystem
C. Cruz, A.M.F. Bio, A. Jullioti, A. Tavares, T. Dias, M.A. Martins-Loução

In Mediterranean-type ecosystems ammonium is present in the soil throughout the year and its concentration is dependent on plant cover.

424 Bidirectional ammonia exchange above a mixed coniferous forest
J. Neirynck, R. Ceulemans

Both stomatal emissions as well as ammonia evaporation from saturated leaves contributed to the canopy emissions from a nitrogen-saturated forest.

439 Airborne ammonia and ammonium within the Northern Adriatic area, Croatia
A. Alebic-Juretic

Airborne ammonia and ammonium for unknown reasons do not follow the reduction in emission levels.

448 Trends of nitrogen in air and precipitation: Model results and observations at EMEP sites in Europe, 1980–2003
H. Fagerli, W. Aas

Ammonium and nitrate in precipitation have declined, whilst the sum of nitrate and nitric acid in air remains at the same level.

462 Genotypic variation of nitrogen use efficiency in Indian mustard
A. Ahmad, I. Khan, Y.P. Abrol, M. Iqbal

Nitrogen efficient crop plants may help in reducing environmental contamination of nitrate without any penalty on seed yield.

467 Atmospheric NH_3 as plant nutrient: A case study with *Brassica oleracea*
A. Castro, I. Stulen, L.J. De Kok

Atmospheric NH_3 can serve as sole N source for Brassica oleracea, but does not change root biomass allocation in nitrate-deprived plants.

473 Low cost measurements of nitrogen and sulphur dry deposition velocities at a semi-alpine site: Gradient measurements and a comparison with deposition model estimates
L.R. Hole, S.H. Brunner, J.E. Hanssen, L. Zhang

Low cost deposition measurement methods need to be tested for different land classes in Europe.

482 Modelling and mapping long-term risks due to reactive nitrogen effects: An overview of LRTAP convention activities
T. Spranger, J.-P. Hettelingh, J. Slootweg, M. Posch

Sustainable targets for European air pollution abatement policy are defined using critical loads in an effects-based approach.

488 Emissions of gaseous nitrogen species from manure management: A new approach
U. Dämmgen, N.J. Hutchings

A cow is a cow. There is no distinction between an IPCC and a UN ECE cow!

498 Estimation of nitrogen balance between the atmosphere and Lake Balaton and a semi natural grassland in Hungary
Sz. Kugler, L. Horváth, A. Machon

The paper summarises the results of works to determine the N-flux between atmosphere and terrestrial/aquatic ecosystems in Hungary.

504 Contents of Volume 154